

4,4',6,6'-Tetrabromo-2,2'-[(*E,E*)-ethane-1,2-diylbis(nitrilomethanylylidene)]diphenol

Hadi Kargar,^a Reza Kia,^{b*‡} Amir Adabi Ardakani^a and Muhammad Nawaz Tahir^{c*}

^aDepartment of Chemistry, Payame Noor University, PO Box 19395-3697 Tehran, I. R. of IRAN, ^bDepartment of Chemistry, Science and Research Branch, Islamic Azad University, Tehran, Iran, and ^cDepartment of Physics, University of Sargodha, Punjab, Pakistan
Correspondence e-mail: zsrkk@yahoo.com, dmntahir_uos@yahoo.com


Received 28 June 2012; accepted 30 June 2012

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}—\text{C}) = 0.010$ Å; R factor = 0.069; wR factor = 0.202; data-to-parameter ratio = 18.2.

The asymmetric unit of the title compound, $\text{C}_{16}\text{H}_{12}\text{Br}_4\text{N}_2\text{O}_2$, comprises half of a potential tetradentate Schiff base ligand. The whole molecule is generated by an inversion center located in the middle of the C—C bond of the ethylene segment. There are intramolecular O—H...N hydrogen bonds making $S(6)$ ring motifs. In the crystal, no significant intermolecular interactions are observed.

Related literature

For standard values of bond lengths, see: Allen *et al.* (1987). For details of hydrogen-bond motifs, see: Bernstein *et al.* (1995). For the crystal structure of a similar compound, see: Kia *et al.* (2012).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{12}\text{Br}_4\text{N}_2\text{O}_2$
 $M_r = 583.92$
Monoclinic, $P2_1/c$
 $a = 12.723$ (3) Å
 $b = 10.291$ (2) Å
 $c = 6.9428$ (18) Å
 $\beta = 97.046$ (15)°
 $V = 902.2$ (4) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 8.93$ mm⁻¹
 $T = 291$ K
 $0.21 \times 0.14 \times 0.08$ mm

Data collection

Bruker SMART APEXII CCD
area-detector diffractometer
Absorption correction: multi-scan
(SADABS; Bruker, 2005)
 $T_{\min} = 0.256$, $T_{\max} = 0.535$
6730 measured reflections
1981 independent reflections
1086 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.075$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.069$
 $wR(F^2) = 0.202$
 $S = 0.98$
1981 reflections
109 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 1.40$ e Å⁻³
 $\Delta\rho_{\min} = -0.90$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1—H1 \cdots N1	0.82	1.83	2.573 (7)	151

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HK and AAA thank PNU for financial support. MNT thanks GC University of Sargodha, Pakistan, for research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2470).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Kia, R., Kargar, H., Adabi Ardakani, A. & Tahir, M. N. (2012). *Acta Cryst.* **E68**, o2242–o2243.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

‡ Present address: Structural Dynamics of (Bio)Chemical Systems, Max Planck Institute for Biophysical Chemistry, Am Fassberg 11, 37077 Göttingen, Germany.

supplementary materials

Acta Cryst. (2012). E68, o2348 [doi:10.1107/S1600536812029832]

4,4',6,6'-Tetrabromo-2,2'-(*E,E*)-ethane-1,2-diylbis(nitrilomethanylylidene)]diphenol

Hadi Kargar, Reza Kia, Amir Adabi Ardakani and Muhammad Nawaz Tahir

Comment

In continuation of our work on the crystal structure analyses of Schiff base ligands (Kargar *et al.*, (2011); Kia *et al.*, (2010), we synthesized the title compound and report herein on its crystal structure.

The asymmetric unit of the title compound, Fig. 1, comprises half of a potentially tetradentate Schiff base ligand. The molecule is located about an inversion center, located in the middle of the C8—C8A bond of the ethylene segment. The bond lengths (Allen *et al.*, 1987) and angles are within the normal ranges and are comparable to those reported for a similar compound (Kia *et al.*, 2012). The intramolecular O—H \cdots N hydrogen bonds (Table 1) make *S*(6) ring motifs (Bernstein *et al.*, 1995).

In the crystal, there are no significant intermolecular interactions present.

Experimental

The title compound was synthesized by adding 3,5-dibromosalicylaldehyde (2 mmol) to a solution of ethylenediamine (1 mmol) in ethanol (30 ml). The mixture was refluxed with stirring for 30 min. The resultant solution was filtered. Yellow single crystals of the title compound suitable for *X*-ray structure determination were recrystallized from ethanol by slow evaporation of the solvents at room temperature over several days.

Refinement

The O-bound H atom was located in a difference Fourier map and constrained to refine on the parent O atom with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. The C-bound H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.93 and 0.97 Å for CH and CH₂ H-atoms, respectively, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Computing details

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).




Figure 1

The molecular structure of the title compound, showing 40% probability displacement ellipsoids and the atomic numbering [symmetry code for suffix A: $-x + 1, -y, -z + 1$]. The intramolecular O-H \cdots N hydrogen bonds are shown as dashed lines.

4,4',6,6'-Tetrabromo-2,2'-(*E,E*)-ethane-1,2-diylbis(nitrilomethanylylidene)diphenol

Crystal data

$C_{16}H_{12}Br_4N_2O_2$
 $M_r = 583.92$
 Monoclinic, $P2_1/c$
 Hall symbol: $-P\ 2_1/c$
 $a = 12.723\ (3)\ \text{\AA}$
 $b = 10.291\ (2)\ \text{\AA}$
 $c = 6.9428\ (18)\ \text{\AA}$
 $\beta = 97.046\ (15)^\circ$
 $V = 902.2\ (4)\ \text{\AA}^3$
 $Z = 2$

$F(000) = 556$
 $D_x = 2.150\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 1563 reflections
 $\theta = 2.5\text{--}27.4^\circ$
 $\mu = 8.93\ \text{mm}^{-1}$
 $T = 291\ \text{K}$
 Block, pale-yellow
 $0.21 \times 0.14 \times 0.08\ \text{mm}$

Data collection

Bruker SMART APEXII CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.256, T_{\max} = 0.535$

6730 measured reflections
 1981 independent reflections
 1086 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.075$
 $\theta_{\max} = 27.1^\circ, \theta_{\min} = 1.6^\circ$
 $h = -16 \rightarrow 13$
 $k = -13 \rightarrow 12$
 $l = -8 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.069$
 $wR(F^2) = 0.202$
 $S = 0.98$
 1981 reflections
 109 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1116P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 1.40\ \text{e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.90\ \text{e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.30804 (8)	0.64680 (8)	0.49803 (17)	0.0684 (4)
Br2	−0.05455 (7)	0.36429 (9)	0.18237 (15)	0.0615 (4)
O1	0.4014 (4)	0.3794 (5)	0.5301 (8)	0.0514 (14)
H1	0.4198	0.3042	0.5527	0.077*
N1	0.3995 (5)	0.1295 (5)	0.5201 (10)	0.0456 (17)
C1	0.3005 (5)	0.3720 (7)	0.4544 (11)	0.0373 (17)
C2	0.2407 (6)	0.4862 (7)	0.4249 (11)	0.0408 (18)
C3	0.1366 (6)	0.4860 (7)	0.3445 (11)	0.0443 (19)
H3A	0.0991	0.5633	0.3236	0.053*
C4	0.0901 (6)	0.3697 (7)	0.2964 (12)	0.0417 (18)
C5	0.1429 (5)	0.2529 (7)	0.3262 (10)	0.0399 (18)
H5A	0.1087	0.1748	0.2926	0.048*
C6	0.2490 (6)	0.2542 (7)	0.4080 (11)	0.0398 (18)
C7	0.3031 (6)	0.1305 (8)	0.4442 (11)	0.044 (2)
H7A	0.2677	0.0529	0.4124	0.053*
C8	0.4522 (6)	0.0026 (8)	0.5551 (13)	0.053 (2)
H8A	0.4034	−0.0668	0.5118	0.063*
H8B	0.4743	−0.0087	0.6928	0.063*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0479 (6)	0.0445 (6)	0.1087 (9)	−0.0044 (4)	−0.0059 (5)	−0.0085 (5)
Br2	0.0317 (5)	0.0595 (7)	0.0885 (8)	0.0053 (3)	−0.0116 (4)	−0.0005 (5)
O1	0.032 (3)	0.051 (3)	0.067 (4)	0.002 (2)	−0.009 (3)	−0.001 (3)
N1	0.040 (4)	0.041 (4)	0.055 (4)	0.014 (3)	0.006 (3)	0.003 (3)
C1	0.022 (4)	0.046 (5)	0.043 (4)	0.004 (3)	0.004 (3)	−0.001 (3)
C2	0.032 (4)	0.043 (4)	0.048 (5)	0.002 (3)	0.002 (3)	−0.004 (4)
C3	0.035 (4)	0.040 (5)	0.055 (5)	0.011 (3)	−0.003 (3)	−0.001 (4)
C4	0.025 (4)	0.046 (5)	0.054 (5)	0.003 (3)	0.002 (3)	−0.001 (4)
C5	0.033 (4)	0.046 (5)	0.040 (4)	−0.004 (3)	0.000 (3)	0.004 (3)
C6	0.038 (4)	0.041 (4)	0.040 (4)	0.007 (3)	0.003 (3)	−0.001 (3)
C7	0.038 (5)	0.052 (5)	0.042 (4)	0.008 (3)	0.003 (3)	−0.003 (4)
C8	0.031 (4)	0.052 (5)	0.074 (6)	0.017 (4)	0.004 (4)	0.013 (4)

Geometric parameters (Å, °)

Br1—C2	1.902 (7)	C3—H3A	0.9300
Br2—C4	1.913 (8)	C4—C5	1.381 (10)
O1—C1	1.328 (9)	C5—C6	1.399 (10)
O1—H1	0.8184	C5—H5A	0.9300
N1—C7	1.274 (10)	C6—C7	1.455 (10)
N1—C8	1.474 (9)	C7—H7A	0.9300
C1—C6	1.396 (10)	C8—C8 ⁱ	1.515 (15)
C1—C2	1.401 (10)	C8—H8A	0.9700
C2—C3	1.372 (10)	C8—H8B	0.9700
C3—C4	1.359 (10)		
C1—O1—H1	105.2	C4—C5—H5A	120.7
C7—N1—C8	118.1 (7)	C6—C5—H5A	120.7
O1—C1—C6	123.0 (6)	C1—C6—C5	120.3 (7)
O1—C1—C2	119.4 (6)	C1—C6—C7	121.4 (7)
C6—C1—C2	117.6 (7)	C5—C6—C7	118.3 (7)
C3—C2—C1	122.5 (7)	N1—C7—C6	119.3 (7)
C3—C2—Br1	119.3 (5)	N1—C7—H7A	120.4
C1—C2—Br1	118.1 (6)	C6—C7—H7A	120.4
C4—C3—C2	118.1 (7)	N1—C8—C8 ⁱ	109.0 (8)
C4—C3—H3A	120.9	N1—C8—H8A	109.9
C2—C3—H3A	120.9	C8 ⁱ —C8—H8A	109.9
C3—C4—C5	122.7 (7)	N1—C8—H8B	109.9
C3—C4—Br2	119.7 (5)	C8 ⁱ —C8—H8B	109.9
C5—C4—Br2	117.5 (6)	H8A—C8—H8B	108.3
C4—C5—C6	118.7 (7)		
O1—C1—C2—C3	179.0 (7)	O1—C1—C6—C5	−179.2 (7)
C6—C1—C2—C3	−3.2 (12)	C2—C1—C6—C5	3.1 (11)
O1—C1—C2—Br1	−0.6 (10)	O1—C1—C6—C7	1.3 (12)
C6—C1—C2—Br1	177.2 (5)	C2—C1—C6—C7	−176.4 (7)
C1—C2—C3—C4	1.5 (12)	C4—C5—C6—C1	−1.3 (11)
Br1—C2—C3—C4	−179.0 (6)	C4—C5—C6—C7	178.2 (7)
C2—C3—C4—C5	0.4 (13)	C8—N1—C7—C6	180.0 (7)
C2—C3—C4—Br2	−179.6 (6)	C1—C6—C7—N1	0.1 (12)
C3—C4—C5—C6	−0.5 (12)	C5—C6—C7—N1	−179.5 (7)
Br2—C4—C5—C6	179.5 (6)	C7—N1—C8—C8 ⁱ	121.0 (10)

Symmetry code: (i) $-x+1, -y, -z+1$.*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots N1	0.82	1.83	2.573 (7)	151